

RELATION BETWEEN ACTION AND ENERGY

Link to: physicspages home page.


To leave a comment or report an error, please use the auxiliary blog.

References: Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Section 2.8; Exercises 2.8.6 - 2.8.7.

Here we'll examine an interesting relation between the action S and the total energy of a system, as given by the Hamiltonian H . Suppose a single particle moving in one dimension follows a classical path given by $x_{cl}(t)$, and moves from an initial position at time t_i of $x_{cl}(t_i) = x_i$ to a final position at time t_f of $x_{cl}(t_f) = x_f$. The action S_{cl} of this classical path is given by the integral of the Lagrangian

$$(1) \quad S_{cl} = \int_{t_i}^{t_f} L(x, \dot{x}) dt$$

What can we say about the rate of change of the action with respect to the final time t_f ? That is, we want to calculate $\partial S_{cl} / \partial t_f$, where all other parameters t_i, x_i and x_f are held constant. The situation can be illustrated as shown:



Since the only thing that is changing is t_f , the particle starts at the same initial time (which we've taken to be $t_i = 0$ in the diagram) and moves to the same location x_f , but at a different time (in the diagram, later time). This

means that the particle must follow a different path, possibly over its entire trajectory. This path, which we'll call $x(t)$, is related to the original path $x_{cl}(t)$ by perturbing the original path by an amount $\eta(t)$:

$$(2) \quad x(t) = x_{cl}(t) + \eta(t)$$

In the diagram, the original path x_{cl} is shown in red and the perturbed path x in blue. The amount η is seen to be the vertical distance between these two curves at each time, and in the case of the paths shown in the diagram, $\eta(t) < 0$.

The difference in the action between the two paths is due to two contributions: first, there is the contribution due to the extra time, from t_f to $t_f + \Delta t$, that the particle takes to complete its path. Second, there is the difference in the two actions over the path from t_i to t_f . The first contribution is entirely new and, for an infinitesimal extra time Δt , it is given by

$$(3) \quad \delta S_1 = L(t_f) \Delta t$$

where $L(t_f)$ is the Lagrangian evaluated at time t_f . The other contribution can be obtained by varying the action over the path from $t_i = 0$ to t_f :

$$(4) \quad \delta S_2 = \int_0^{t_f} \delta L dt$$

Since L depends on x and \dot{x} , we have

$$(5) \quad \delta L = \frac{\partial L}{\partial x} \delta x + \frac{\partial L}{\partial \dot{x}} \delta \dot{x}$$

For infinitesimally different trajectories, we can see from the diagram above that $\delta x = \eta(t)$ at each point on the curve, so $\delta \dot{x} = \dot{\eta}(t)$, so we get

$$(6) \quad \delta S_2 = \int_0^{t_f} \left[\frac{\partial L}{\partial x} \eta(t) + \frac{\partial L}{\partial \dot{x}} \dot{\eta}(t) \right] dt$$

$$(7) \quad = \int_0^{t_f} \left[-\frac{d}{dt} \frac{\partial L}{\partial \dot{x}} + \frac{\partial L}{\partial x} \right] \eta(t) dt + \int_0^{t_f} \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \eta(t) \right) dt$$

$$(8) \quad = 0 + \left. \frac{\partial L}{\partial \dot{x}} \eta(t) \right|_{t_f}$$

In these equations, the derivatives of L are evaluated on the original curve x_{cl} . To verify the second line, use the product rule on the second integrand

and cancel terms to get the first line. The second term in the last is evaluated at $t = t_f$ only since we're assuming that $\eta(0) = 0$.

The quantity in brackets in the first integral is zero, because of the Euler-Lagrange equations which are valid on the original curve x_{cl} :

$$(9) \quad \frac{d}{dt} \frac{\partial L}{\partial \dot{x}} - \frac{\partial L}{\partial x} = 0$$

Putting everything together, we get for the total variation in the action:

$$(10) \quad \delta S_{cl} = \delta S_1 + \delta S_2$$

$$(11) \quad = \left[\frac{\partial L}{\partial \dot{x}} \eta(t) + L \Delta t \right]_{t_f}$$

Looking at the diagram above, the slope of the blue curve $x(t_f)$ at the time t_f is given by

$$(12) \quad \dot{x}(t_f) = \frac{|\eta(t_f)|}{\Delta t}$$

From the definition 2 of η we see that $\eta(t_f) < 0$, so

$$(13) \quad \eta(t_f) = -\dot{x}(t_f) \Delta t$$

This gives the final equation for the variation of the action:

$$(14) \quad \delta S_{cl} = \left[-\frac{\partial L}{\partial \dot{x}} \dot{x} + L \right]_{t_f} \Delta t$$

$$(15) \quad = (-p\dot{x} + L) \Delta t$$


$$(16) \quad = -H \Delta t$$

where the second line follows from the definition of the canonical momentum $p = \partial L / \partial \dot{x}$.

The required derivative is

$$(17) \quad \boxed{\frac{\partial S_{cl}}{\partial t_f} = -H(t_f)}$$

Using a similar technique, we can work out $\partial S_{cl} / \partial x_f$. In this case, the situation is as shown in this diagram:



The two trajectories now take the same time, but in the modified trajectory, the particle moves a distance Δx further. Since both paths take the same time, there is no extra contribution $L\Delta t$. In this case $\eta(t) > 0$, since the new (blue) curve $x(t)$ is above the old (red) one $x_{cl}(t)$. The derivation is the same as above up to 8, and the total variation in the action is now

$$(18) \quad \delta S_{cl} = \left. \frac{\partial L}{\partial \dot{x}} \eta(t) \right|_{t_f}$$

At $t = t_f$, $\eta(t_f) = \Delta x$, so we get

$$(19) \quad \delta S_{cl} = \left. \frac{\partial L}{\partial \dot{x}} \right|_{t_f} \Delta x$$

$$(20) \quad \frac{\partial S_{cl}}{\partial x_f} = \left. \frac{\partial L}{\partial \dot{x}} \right|_{t_f} = p(t_f)$$

Example. We can verify 17 for the case of the one-dimensional harmonic oscillator. The general solution for the position is given by

$$(21) \quad x(t) = A \cos \omega t + B \sin \omega t$$

$$(22) \quad \dot{x}(t) = -A\omega \sin \omega t + B\omega \cos \omega t$$

The total energy is given by

$$(23) \quad E = \frac{1}{2}m\dot{x}^2 + \frac{1}{2}m\omega^2 x^2$$

$$(24) \quad = \frac{m}{2} \left((-A\omega \sin \omega t + B\omega \cos \omega t)^2 + \omega^2 (A \cos \omega t + B \sin \omega t)^2 \right)$$

$$(25) \quad = \frac{m\omega^2}{2} (A^2 + B^2)$$

where we just multiplied out the second line, cancelled terms and used $\cos^2 x + \sin^2 x = 1$.

To get the action, we need the Lagrangian:

$$(26)$$

$$L = T - V$$

$$(27)$$

$$= \frac{1}{2}m\dot{x}^2 - \frac{1}{2}m\omega^2 x^2$$

$$(28)$$

$$= \frac{m}{2} \left((-A\omega \sin \omega t + B\omega \cos \omega t)^2 - \omega^2 (A \cos \omega t + B \sin \omega t)^2 \right)$$

$$(29)$$

$$= \frac{m\omega^2}{2} [A^2 (\sin^2 \omega t - \cos^2 \omega t) + B^2 (\cos^2 \omega t - \sin^2 \omega t) - 4AB \sin \omega t \cos \omega t]$$

$$(30)$$

$$= \frac{m\omega^2}{2} ((B^2 - A^2) \cos 2\omega t - 2AB \sin 2\omega t)$$

The action for a trajectory from $t = 0$ to $t = T$ is then

$$(31) \quad S = \int_0^T L dt$$

$$(32) \quad = \frac{m\omega}{4} [(B^2 - A^2) \sin 2\omega t + 2AB \cos 2\omega t]_0^T$$

$$(33) \quad = \frac{m\omega}{4} [(B^2 - A^2) \sin 2\omega T + 2AB (\cos 2\omega T - 1)]$$

$$(34) \quad = \frac{m\omega}{2} [(B^2 - A^2) \sin \omega T \cos \omega T + AB (\cos^2 \omega T - \sin^2 \omega T - 1)]$$

$$(35) \quad = \frac{m\omega}{2} [(B^2 - A^2) \sin \omega T \cos \omega T - 2AB \sin^2 \omega T]$$

To proceed further, we need to specify A and B , since these depend on the boundary conditions (that is, on where we require the mass to be at $t = 0$ and $t = T$). If we require $x(0) = x_1$ and $x(T) = x_2$, then

$$(36) \quad A = x_1$$

$$(37) \quad x_1 \cos \omega T + B \sin \omega T = x_2$$

$$(38) \quad B = \frac{x_2 - x_1 \cos \omega T}{\sin \omega T}$$

Plugging these into 25 gives the energy as

$$(39) \quad E = \frac{m\omega^2}{2} \left(x_1^2 + \left(\frac{x_2 - x_1 \cos \omega T}{\sin \omega T} \right)^2 \right)$$

$$(40) \quad = \frac{m\omega^2}{2 \sin^2 \omega T} (x_1^2 + x_2^2 - 2x_1x_2 \cos \omega T)$$

Plugging A and B into 35, we get (using $c \equiv \cos \omega T$ and $s \equiv \sin \omega T$, so that $s^2 + c^2 = 1$):

$$(41) \quad S = \frac{m\omega}{2s} \left[(x_2 - x_1c)^2 c - x_1s^2c - 2x_1s^2(x_2 - x_1c) \right]$$

$$(42) \quad = \frac{m\omega}{2s} \left[(x_2^2 - 2x_1x_2c + x_1^2c^2) c - x_1^2s^2c - 2x_1x_2s^2 + 2x_1s^2c \right]$$

$$(43) \quad = \frac{m\omega}{2s} \left[(x_1^2 + x_2^2) c - 2x_1x_2 \right]$$

$$(44) \quad = \frac{m\omega}{2 \sin \omega T} \left[(x_1^2 + x_2^2) \cos \omega T - 2x_1x_2 \right]$$

Taking the derivative, we get

$$(45) \quad \frac{\partial S}{\partial T} = \frac{m\omega}{2s^2} \left[-\omega (x_1^2 + x_2^2) s^2 - ((x_1^2 + x_2^2) c - 2x_1x_2) \omega c \right]$$

$$(46) \quad = \frac{m\omega^2}{2s^2} \left[-(x_1^2 + x_2^2) + 2x_1x_2c \right]$$

$$(47) \quad = -\frac{m\omega^2}{2 \sin^2 \omega T} (x_1^2 + x_2^2 - 2x_1x_2 \cos \omega T)$$

$$(48) \quad = -E$$

Thus the result is verified for the harmonic oscillator.

PINGBACKS

Pingback: Path integrals for special potentials; use of classical action