

POSTULATES OF QUANTUM MECHANICS: STATES AND MEASUREMENTS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

References: Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Sections 4.1 - 4.2; Exercise 4.2.1.

Although we've covered the basics of nonrelativistic quantum mechanics before, the approach taken by Shankar in his Chapter 4 provides a new way of looking at it, so it's worth a summary.

Quantum mechanics is based on four postulates, the first three of which describe the quantum state at a fixed instant in time, and the fourth which describes its time evolution via the Schrödinger equation. We'll summarize the first three postulates here, and compare each with its classical analogue.

First, in classical mechanics, the path of a particle is, in the Hamiltonian formalism, described by specifying its position $x(t)$ and momentum $p(t)$ as functions of time. Both the position and momentum are specified precisely at all times. In quantum mechanics, the state of a particle is specified by a vector (ket) $|\psi(t)\rangle$ in a Hilbert space.

Second, in classical mechanics, any dynamical variable ω is a function of the two phase-space coordinates x and p : $\omega = \omega(x, p)$. In quantum mechanics, the spatial coordinate x is replaced by the Hermitian operator X and the momentum p is replaced by the differential operator $P = \hbar K$ which we discussed earlier. The matrix elements of X and P in position space are

$$(0.1) \quad \langle x|X|x'\rangle = x\delta(x-x')$$

$$(0.2) \quad \langle x|P|x'\rangle = -i\hbar\delta'(x-x')$$

The classical dynamical variable $\omega(x, p)$ becomes a Hermitian operator $\Omega(X, P)$, where x and p in $\omega(x, p)$ are replaced by their corresponding operators X and P .

The third postulate states how measurements work in quantum mechanics. In classical mechanics, it is assumed that (in principle) any dynamical variable ω may be measured with arbitrary precision without changing the state of the particle. In quantum mechanics, if we wish to measure the value of a variable represented by the operator Ω , we must determine the eigenvalues ω_i and corresponding eigenvectors $|\omega_i\rangle$ of Ω , then express the state $|\psi\rangle$ as a linear combination of the $|\omega_i\rangle$. Then the best we can do is to state that

the particular eigenvalue ω_i will be measured with probability $|\langle \omega_i | \psi \rangle|^2$. After the measurement, the state $|\psi\rangle$ 'collapses' to become the state $|\omega_i\rangle$. The only possible outcomes of a measurement of Ω are its eigenvalues; no intermediate values are possible.

To illustrate these postulates, suppose we have the following three operators on a complex 3-d Hilbert space (essentially these are the spin-1 operators without the \hbar)

$$(0.3) \quad L_x = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$(0.4) \quad L_y = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{bmatrix}$$

$$(0.5) \quad L_z = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

Since L_z is diagonal, its eigenvalues can be read off from the diagonal elements as $0, \pm 1$, so these are the possible values of L_z that could be obtained in a measurement. Also because L_z is diagonal, its eigenvectors are

$$(0.6) \quad |L_z = +1\rangle = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$(0.7) \quad |L_z = 0\rangle = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

$$(0.8) \quad |L_z = -1\rangle = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Suppose we start with the state $|L_z = +1\rangle$ in which $L_z = +1$, and we want to measure L_x in this state. To find the expectation values $\langle L_x \rangle$ and $\langle L_x^2 \rangle$ in this state, we calculate

$$\begin{aligned}
 (0.9) \quad \langle L_x \rangle &= \langle L_z = +1 | L_x | L_z = +1 \rangle \\
 (0.10) \quad &= [1 \ 0 \ 0] \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \\
 (0.11) \quad &= \frac{1}{\sqrt{2}} [1 \ 0 \ 0] \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \\
 (0.12) \quad &= 0
 \end{aligned}$$

To get $\langle L_x^2 \rangle$ we first find the operator

$$(0.13) \quad L_x^2 = \frac{1}{2} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

Now we have

$$\begin{aligned}
 (0.14) \quad \langle L_x^2 \rangle &= [1 \ 0 \ 0] \frac{1}{2} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \\
 (0.15) \quad &= \frac{1}{2} [1 \ 0 \ 0] \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \\
 (0.16) \quad &= \frac{1}{2}
 \end{aligned}$$

The uncertainty, or variance, is

$$(0.17) \quad \Delta L_x = \sqrt{\langle L_x^2 \rangle - \langle L_x \rangle^2} = \frac{1}{\sqrt{2}}$$

To find the possible values of L_x and their probabilities, we need to find the eigenvalues and eigenvectors of L_x , which we can do in the L_z basis, since this basis is given by the three vectors in 0.6. The eigenvalues are found in the usual way from the determinant:

$$(0.18) \quad \begin{vmatrix} -\lambda & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & -\lambda & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & -\lambda \end{vmatrix} = -\lambda \left(\lambda^2 - \frac{1}{2} \right) - \frac{1}{\sqrt{2}} \left(\frac{-\lambda}{\sqrt{2}} \right)$$

$$(0.19) \quad = -\lambda^3 + \lambda = 0$$

$$(0.20) \quad \lambda = 0, \pm 1$$

The eigenvectors can be found in the usual way, by solving

$$(0.21) \quad (L_x - \lambda I) |L_x = \lambda\rangle = 0$$

where the ket takes on the three possible values of λ successively. We let

$$(0.22) \quad |L_x = \lambda\rangle = \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

For $\lambda = +1$ we have

$$(0.23) \quad -a + \frac{b}{\sqrt{2}} = 0$$

$$(0.24) \quad \frac{1}{\sqrt{2}} (a - \sqrt{2}b + c) = 0$$

$$(0.25) \quad \frac{b}{\sqrt{2}} - c = 0$$

Only two of these three equations are independent, so we can set $a = 1$ and solve for b and c to get

$$(0.26) \quad a = 1$$

$$(0.27) \quad b = \sqrt{2}$$

$$(0.28) \quad c = 1$$

Normalizing the eigenvector gives

$$(0.29) \quad |L_x = +1\rangle = \frac{1}{2} \begin{bmatrix} 1 \\ \sqrt{2} \\ 1 \end{bmatrix}$$

The other two eigenvectors can be found the same way, with the result

$$(0.30) \quad |L_x = 0\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$(0.31) \quad |L_x = -1\rangle = \frac{1}{2} \begin{bmatrix} 1 \\ -\sqrt{2} \\ 1 \end{bmatrix}$$

Note that these eigenvectors are orthonormal.

Now that we have the eigenvectors of L_x we can answer the following question. If we start with the state $|L_z = -1\rangle$ and measure L_x , what are the possible outcomes and the probability of each?

First, we need to express $|L_z = -1\rangle$ in terms of the eigenvectors of L_x which we can do by solving three simultaneous linear equations, and we find

$$(0.32) \quad |L_z = -1\rangle = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \frac{1}{2} (|L_x = +1\rangle + |L_x = -1\rangle) - \frac{1}{\sqrt{2}} |L_x = 0\rangle$$

(You can verify this by direct substitution.) Thus all 3 possible values of L_x can result from a measurement, and the probability of each is

$$(0.33) \quad P(L_x = +1) = |\langle L_x = +1 | L_z = -1 \rangle|^2$$

$$(0.34) \quad = \left(\frac{1}{2} [1 \quad \sqrt{2} \quad 1] \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)^2$$

$$(0.35) \quad = \frac{1}{4}$$

$$(0.36) \quad P(L_x = 0) = |\langle L_x = 0 | L_z = -1 \rangle|^2$$

$$(0.37) \quad = \left(\frac{1}{\sqrt{2}} [1 \quad 0 \quad -1] \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)^2$$

$$(0.38) \quad = \frac{1}{2}$$

$$(0.39) \quad P(L_x = -1) = |\langle L_x = -1 | L_z = -1 \rangle|^2$$

$$(0.40) \quad = \left(\frac{1}{2} [1 \quad -\sqrt{2} \quad 1] \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)^2$$

$$(0.41) \quad = \frac{1}{4}$$

Now suppose we start with the state, written in the L_z basis:

$$(0.42) \quad |\psi\rangle = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

We take a measurement of L_z^2 and obtain $+1$. The operator L_z^2 is given by squaring 0.5:

$$(0.43) \quad L_z^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

This has a degenerate eigenvalue $\lambda = +1$, so the most we can say about the state $|\psi\rangle$ after the measurement is that it is projected onto the subspace

$a \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + b \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$. That is, the state after the measurement is given by

$$(0.44) \quad |\psi\rangle_{after} = \mathbb{P}_{L_z=\pm 1} |\psi\rangle_{before}$$

$$(0.45) \quad = [|L_z = +1\rangle \langle L_z = +1| + |L_z = -1\rangle \langle L_z = -1|] \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$(0.46) \quad = \left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} [1 \ 0 \ 0] + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} [0 \ 0 \ 1] \right) \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$(0.47) \quad = \begin{bmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

We can normalize this state to get

$$(0.48) \quad |\psi\rangle_{after} = \frac{2}{\sqrt{3}} \begin{bmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Thus if we measure L_z immediately after the measurement of L_z^2 above, we get $L_z = +1$ with probability $\frac{1}{3}$ and $L_z = -1$ with probability $\frac{2}{3}$.

Finally, suppose we have a state $|\psi\rangle$ with the probabilities of measurements of L_z given as $P(L_z = 1) = \frac{1}{4}$, $P(L_z = 0) = \frac{1}{2}$ and $P(L_z = -1) = \frac{1}{4}$. Since these probabilities are given by $|\langle L_z = \lambda | \psi \rangle|^2$ for each of the three possible values of λ , and the vectors $|L_z = \lambda\rangle$ are orthonormal, the most general form for $|\psi\rangle$ is

$$(0.49) \quad |\psi\rangle = \frac{e^{i\delta_1}}{2} |L_z = 1\rangle + \frac{e^{i\delta_2}}{\sqrt{2}} |L_z = 0\rangle + \frac{e^{i\delta_3}}{2} |L_z = -1\rangle$$

where the δ_i are real numbers. For example

$$(0.50) \quad |\langle L_z = 1 | \psi \rangle|^2 = \left| \frac{e^{i\delta_1}}{2} \right|^2 = \frac{1}{4}$$

While the presence of a phase factor in a solitary state doesn't affect the physics of that state, if we have a sum of states, each with its own (different) phase factor, we can't ignore these phase factors. For example, if we measure L_x in this state and want the probability that $L_x = 0$, we have, using 0.30

(0.51)

$$P(L_x = 0) = |\langle L_x = 0 | \psi \rangle|^2$$

(0.52)

$$= \left| \frac{1}{\sqrt{2}} [1 \ 0 \ -1] \left(\frac{e^{i\delta_1}}{2} |L_z = 1\rangle + \frac{e^{i\delta_2}}{\sqrt{2}} |L_z = 0\rangle + \frac{e^{i\delta_3}}{2} |L_z = -1\rangle \right) \right|^2$$

(0.53)

$$= \left| \frac{1}{\sqrt{2}} [1 \ 0 \ -1] \left(\frac{e^{i\delta_1}}{2} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \frac{e^{i\delta_2}}{\sqrt{2}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + \frac{e^{i\delta_3}}{2} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right) \right|^2$$

(0.54)

$$= \frac{1}{8} |e^{i\delta_1} - e^{i\delta_3}|^2$$

(0.55)

$$= \frac{1}{8} |1 - e^{i(\delta_3 - \delta_1)}|^2$$

The last line will have a different result for different values of the phase factors δ_1 and δ_3 , so they can't be ignored.

PINGBACKS

Pingback: Postulates of quantum mechanics: momentum

Pingback: Postulates of quantum mechanics: Schrödinger equation and propagators

Pingback: Poisson brackets to commutators: classical to quantum