

## POSTULATES OF QUANTUM MECHANICS: MOMENTUM

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Sections 4.1 - 4.2; Exercises 4.2.2 - 4.2.3.

One of the postulates of quantum mechanics is that the momentum operator  $P$  in position space is given by

$$(1) \quad \langle x|P|x'\rangle = -i\hbar\delta'(x-x')$$

By using the properties of the derivative of the delta function, we can find the eigenfunctions of  $P$ . We have

$$(2) \quad \langle x|P|\psi\rangle = \int \langle x|P|x'\rangle \langle x'|\psi\rangle dx'$$

$$(3) \quad = -i\hbar \int \delta'(x-x') \langle x'|\psi\rangle dx'$$

$$(4) \quad = -i\hbar \frac{d}{dx} \langle x|\psi\rangle$$

$$(5) \quad = -i\hbar \frac{d\psi(x)}{dx}$$

The eigenvector of  $P$  is  $|p\rangle$  and has the property that

$$(6) \quad P|p\rangle = p|p\rangle$$

If we project this onto position space and use 5 we get

$$(7) \quad \langle x|P|\psi\rangle = p \langle x|p\rangle$$

$$(8) \quad -i\hbar \frac{d\psi_p(x)}{dx} = p\psi_p(x)$$

where

$$(9) \quad \psi_p(x) \equiv \langle x|p\rangle$$

Solving this differential equation and normalizing so that  $\langle p'|p\rangle = \delta(p-p')$  we get

$$(10) \quad \psi_p(x) = \frac{1}{\sqrt{2\pi\hbar}} e^{ipx/\hbar}$$

For an arbitrary wave function  $|\psi\rangle$ , if we know its position-space form, we can find its momentum-space version as follows:

$$(11) \quad \langle p|\psi\rangle = \int \langle p|x\rangle \langle x|\psi\rangle dx$$

$$(12) \quad = \int \psi_p^*(x) \langle x|\psi\rangle dx$$

$$(13) \quad = \frac{1}{\sqrt{2\pi\hbar}} \int e^{-ipx/\hbar} \psi(x) dx$$

This has an interesting consequence if the position-space function  $\psi(x)$  is real. The probability density for finding a particle in a state with momentum  $p$  is  $|\langle p|\psi\rangle|^2$ , which we can write as

$$(14) \quad |\langle p|\psi\rangle|^2 = \langle p|\psi\rangle^* \langle p|\psi\rangle$$

$$(15) \quad = \frac{1}{2\pi\hbar} \int \int e^{ip(x-x')/\hbar} \psi(x) \psi(x') dx dx'$$

$$(16) \quad = \frac{1}{2\pi\hbar} \int \int e^{-ip(x'-x)/\hbar} \psi(x) \psi(x') dx dx'$$

$$(17) \quad = \frac{1}{2\pi\hbar} \int \int e^{-ip(x-x')/\hbar} \psi(x') \psi(x) dx dx'$$

$$(18) \quad = |\langle -p|\psi\rangle|^2$$

In the fourth line, since  $x$  and  $x'$  are dummy integration variables, both of which are integrated over the same range, we can simply swap them without changing anything. Note that the derivation relies on  $\psi(x)$  being real, since if it were complex we would have

$$\begin{aligned}
(19) \quad & |\langle p | \psi \rangle|^2 = \langle p | \psi \rangle^* \langle p | \psi \rangle \\
(20) \quad & = \frac{1}{2\pi\hbar} \int \int e^{ip(x-x')/\hbar} \psi(x) \psi^*(x') dx dx' \\
(21) \quad & = \frac{1}{2\pi\hbar} \int \int e^{-ip(x'-x)/\hbar} \psi(x) \psi^*(x') dx dx' \\
(22) \quad & = \frac{1}{2\pi\hbar} \int \int e^{-ip(x-x')/\hbar} \psi(x') \psi^*(x) dx dx' \\
(23) \quad & \neq |\langle -p | \psi \rangle|^2
\end{aligned}$$

since

$$(24) \quad |\langle -p | \psi \rangle|^2 = \frac{1}{2\pi\hbar} \int \int e^{-ip(x-x')/\hbar} \psi(x) \psi^*(x') dx dx'$$

That is, for  $|\langle -p | \psi \rangle|^2$  the position  $x'$  that is the argument of the  $\psi^*(x')$  factor appears as the positive term  $ipx'$  in the exponential, but in 22 the argument of the complex conjugate wave function is  $x$ , which appears as the negative term  $-ipx$  in the exponential.

Thus for any real wave function, the probability of the particle having momentum  $+p$  is equal to the probability of it having  $-p$ , so for such wave functions, the mean momentum is always  $\langle P \rangle = 0$ .

As another example, suppose we have a wave function  $\psi(x)$  with a mean momentum  $\bar{p}$ , so that

$$(25) \quad \langle \psi | P | \psi \rangle = \bar{p}$$

If we now multiply  $\psi$  by  $e^{ip_0x/\hbar}$  where  $p_0$  is a constant momentum, we can calculate the new mean momentum using 5:

$$\begin{aligned}
(26) \quad & \langle P \rangle = \langle e^{ip_0x/\hbar} \psi | P | e^{ip_0x/\hbar} \psi \rangle \\
(27) \quad & = -i\hbar \int e^{-ip_0x/\hbar} \psi^*(x) \frac{d}{dx} \left( e^{ip_0x/\hbar} \psi(x) \right) dx \\
(28) \quad & = -i\hbar \int e^{-ip_0x/\hbar} \psi^* \left[ \frac{ip_0}{\hbar} e^{ip_0x/\hbar} \psi(x) + e^{ip_0x/\hbar} \frac{d}{dx} \psi(x) \right] dx \\
(29) \quad & = \int p_0 \psi^* \psi dx - i\hbar \int \psi^*(x) \frac{d}{dx} \psi(x) dx \\
(30) \quad & = p_0 + \bar{p}
\end{aligned}$$

The first integral in the fourth line uses the fact that  $p_0$  is constant and  $\psi$  is normalized so that

$$(31) \quad \int \psi^* \psi dx = 1$$

PINGBACKS

Pingback: Free particle in the position basis

Pingback: Propagator for a Gaussian wave packet for the free particle