

CHANGING THE POSITION BASIS WITH A UNITARY TRANSFORMATION

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Section 7.4, Exercise 7.4.9.

The standard representation of the position and momentum operators in the position basis is

$$(0.1) \quad X \rightarrow x$$

$$(0.2) \quad P \rightarrow -i\hbar \frac{d}{dx}$$

It turns out it's possible to modify this definition by adding some arbitrary function of position $f(x)$ to P so we have

$$(0.3) \quad X' \rightarrow x$$

$$(0.4) \quad P' \rightarrow -i\hbar \frac{d}{dx} + f(x)$$

Since any function of x commutes with X , the commutation relations remain unchanged, so we have

$$(0.5) \quad [X', P'] = i\hbar$$

Another way of interpreting this change in operators is by using the unitary transformation of the X basis, in the form

$$(0.6) \quad |x\rangle \rightarrow |\tilde{x}\rangle = e^{ig(X)/\hbar} |x\rangle = e^{ig(x)/\hbar} |x\rangle$$

where

$$(0.7) \quad g(x) \equiv \int^x f(x') dx'$$

The last equality in 0.6 comes from the fact that operating on $|x\rangle$ with any function of the X operator (provided the function can be expanded in a power series) results in multiplying $|x\rangle$ by the same function, but with the operator X replaced by the numeric position value.

To verify this works, we can calculate the matrix elements of the old X and P operators in the new basis. We have

$$(0.8) \quad \langle \tilde{x} | X | \tilde{x}' \rangle = \left\langle x \left| e^{-ig(x)/\hbar} X e^{ig(x')/\hbar} \right| x' \right\rangle$$

At this stage, since the two exponentials are numerical functions and not operators, we can take them outside the bracket to

$$(0.9) \quad \langle \tilde{x} | X | \tilde{x}' \rangle = e^{-ig(x)/\hbar} e^{ig(x')/\hbar} \langle x | X | x' \rangle$$

$$(0.10) \quad = e^{-ig(x)/\hbar} e^{ig(x')/\hbar} x' \delta(x - x')$$

$$(0.11) \quad = x \delta(x - x')$$

The exponentials cancel in the last line since the delta function is non-zero only when $x = x'$.

The above result can also be obtained by inserting a couple of identity operators into 0.8:

$$(0.12)$$

$$\left\langle x \left| e^{-ig(x)/\hbar} X e^{ig(x')/\hbar} \right| x' \right\rangle = \int \int \left\langle x \left| e^{-ig(x)/\hbar} \right| y \right\rangle \langle y | X | z \rangle \left\langle z \left| e^{ig(x')/\hbar} \right| x' \right\rangle dy dz$$

$$(0.13) \quad = \int \int \left\langle x \left| e^{-ig(x)/\hbar} \right| y \right\rangle z \delta(y - z) \left\langle z \left| e^{ig(x')/\hbar} \right| x' \right\rangle dy dz$$

$$(0.14) \quad = \int \left\langle x \left| e^{-ig(x)/\hbar} \right| z \right\rangle z \left\langle z \left| e^{ig(x')/\hbar} \right| x' \right\rangle dz$$

$$(0.15) \quad = \int e^{i[g(x') - g(x)]/\hbar} \langle x | z \rangle z \langle z | x' \rangle dz$$

$$(0.16) \quad = \int e^{i[g(x') - g(x)]/\hbar} \delta(x - z) z \delta(z - x') dz$$

$$(0.17) \quad = e^{i[g(x') - g(x)]/\hbar} x' \delta(x - x')$$

$$(0.18) \quad = x \delta(x - x')$$

The momentum operator works as follows. Using the original definition 0.2 on the modified basis we have

$$(0.19) \quad \langle \tilde{x} | P | \tilde{x}' \rangle = -i\hbar \left\langle x \left| e^{-ig(x)/\hbar} \frac{d}{dx'} e^{ig(x')/\hbar} \right| x' \right\rangle$$

$$(0.20) \quad = -i\hbar \left\langle x \left| e^{-ig(x)/\hbar} \frac{i}{\hbar} e^{ig(x')/\hbar} \frac{dg(x')}{dx'} \right| x' \right\rangle -$$

$$(0.21) \quad i\hbar \left\langle x \left| e^{-ig(x)/\hbar} e^{ig(x')/\hbar} \frac{d}{dx'} \right| x' \right\rangle$$

From 0.7 we have

$$(0.22) \quad \frac{dg(x)}{dx} = \frac{d}{dx} \int^x f(x') dx' = f(x)$$

This gives

$$(0.23) \quad \langle \tilde{x} | P | \tilde{x}' \rangle = \left\langle x \left| e^{i[g(x')-g(x)]/\hbar} \left[f(x') - i\hbar \frac{d}{dx'} \right] \right| x' \right\rangle$$

$$(0.24) \quad = e^{i[g(x')-g(x)]/\hbar} \left[f(x') - i\hbar \frac{d}{dx'} \right] \langle x | x' \rangle$$

$$(0.25) \quad = e^{i[g(x')-g(x)]/\hbar} \left[f(x') - i\hbar \frac{d}{dx'} \right] \delta(x-x')$$

$$(0.26) \quad = \left[f(x) - i\hbar \frac{d}{dx} \right] \delta(x-x')$$

This shows that by a unitary change of X basis 0.6, we transform the position and momentum operators (well, just the momentum operator, really) according to 0.3. We've multiplied the original $|x\rangle$ states by a phase factor which depends on some function $f(x)$. This doesn't change the matrix elements of X , but it does add $f(x)$ to the matrix elements of P . The commonly used definition of P is thus with $f(x) = 0$.